

All Databases Journals PubMed Books Nucleotide Protein Genome Structure OMIM PMC

Search for

Advanced Search

[Limits](#) [Preview/Index](#) [History](#) [Clipboard](#) [Details](#)Display Show Sort By Send to

Alt: 1 Review: 0

1: [Cancer Res. 2006 Feb 15;66\(4\):1949-55.](#) [Links](#)**Identification of the IFITM family as a new molecular marker in human colorectal tumors.****Andreu P, Colnot S, Godard C, Laurent-Puig P, Lamarque D, Kahn A, Perret C, Romagnolo B.**

Institut Cochin, INSERM U567, Centre National de la Recherche Scientifique UMR8104, Université Paris V, France.

We analyzed the expression profiles of intestinal adenomas from a new murine familial adenomatous polyposis model (Apc(delta14/+)) using suppression subtractive hybridization to identify novel diagnostic markers of colorectal carcinogenesis. We identified 18 candidate genes having increased expression levels in the adenoma. Subsequent Northern blotting, real-time reverse transcription-PCR, and in situ hybridization analysis confirmed their induction in beta-catenin-activated epithelial cells of murine adenomas. We showed that most of the genes also have altered expression levels in human colonic adenomas and carcinomas. We focused on the IFITM genes that encode IFN-inducible transmembrane proteins. Serial analyses of gene expression levels revealed high levels of expression in early and late intestinal neoplasm in both mice and humans. Using a conditional mouse model of Apc inactivation and a human colon carcinoma cell line, we showed that IFITM gene expression is rapidly induced after activation of the beta-catenin signaling. Using a large-scale analysis of human tumors, we showed that IFITM gene expression is significantly up-regulated specifically in colorectal tumors and thus may be a useful diagnostic tool in these tumors.

PMID: 16488993 [PubMed - indexed for MEDLINE]

Related articles

Differential gene expression profile reveals deregulation of the cancer-specific beta1 glycoprotein 9 early during colorectal carcinogenesis. [Cancer Res. 2005]

A colorectal cancer expression profile that includes transforming growth factor-beta inhibitor BAMBI predicts metastatic potential. [Cancer Res. 2005]

Beta-catenin/Tcf-4 inhibition after progesterin target gene induction drives differentiation of intestinal tumors. [Cancer Res. 2005]

[Review](#) Wnt signaling inside the nucleus. [Cancer Sci. 2005]

[Review](#) Colorectal cancer and genetic alterations in the Wnt pathway. [Cancer Res. 2005]

» See reviews... | » See all...

Cited by 4 PubMed Central articles

[Review](#) Target Identification for CNS diseases by transcriptional profiling. 2009]

Normal germ line establishment in mice carrying a deletion of the *Wnt1* gene. [Development. 2008]

Up-regulation of NG2 proteoglycan and interferon-induced transmembrane proteins 1 and 3 in mouse astrocytoma: a membrane proteomics approach. [Mol Cell Proteomics. 2006]

» See all...

Recent Activity

[Turn Off](#) [Clear](#)

Identification of the IFITM family
as a new molecular marker in
human colorectal tumors.

IFITM2 (13)

human 1-8D interferon ind... (11)

{human 1-8D} AND and AND ...
(11)

[PubMed](#)

Functional detection of epithelial
cell adhesion molecule specific
cytotoxic T lymphocytes...

[» See more...](#)

Display [AbstractPlus](#) Show [20](#) Sort By [Send to](#)

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)